

Qualitative and quantitative composition

Ivermectin 0.02% w/w (200µg/ml)

Pharmaceutical form

Topical spray.

Pharmacological properties

XENO 200 Spray is designed as a topical spray for use on rabbits, guinea pigs, rodents, ferrets, birds and reptiles. The active ingredient Ivermectin is a substance known as an endectocide because it kills parasites that cause infestation inside the body (endoparasites) as well as those living outside the body, on the skin (ectoparasites). Ivermectin is used in many species for the control of mites, roundworms and lice.

Clinical particulars

Target species

Rabbits, guinea pigs, rodents, ferrets, birds and reptiles.

Indications for use

Prevents and treats common internal and external parasites of rabbits, guinea pigs, rodents, ferrets, birds and reptiles.

Contra-indications

XENO 200 Spray is only for use on rabbits, guinea pigs, ferrets, rodents reptiles and birds. Do not use in animals other than those indicated. Serious reactions including deaths have been reported in dogs (especially Collies, Old English Sheepdogs and related breeds), tortoises and turtles treated with the active substance.

Special precautions for use

If you are uncertain about the condition of your pet, consult a veterinary surgeon. If symptoms persist, consult a veterinary surgeon.

Use during pregnancy, lactation or lay

No data available. Do not use in pregnant and lactating animals.

Interactions

None known.

Amounts to be administered and administration route

In order to apply a correct dosage it is essential to accurately weigh the animal. Your veterinary practice can help you to do this. Using the table below calculate the correct amount of XENO 200 Spray to use and how many pump actuations you should give.

XENO 200 Spray – Dosage Table			
SPECIES	Remarks	Dose rate	AMOUNT TO USE (number of pump actuations)
Rabbits, Guinea pigs and Ferrets		400µg/kg bodyweight	1 pump per 100g bodyweight (0.2ml) 10 pumps per kg bodyweight (2ml)
Small Rodents	Do not use in animals under 100g bodyweight	200µg/kg bodyweight	1 pump per 200g bodyweight (0.2ml)

Reptiles	DO NOT USE IN Tortoises, Turtles, Terrapins, Indigo, Snakes or Skinks	200µg/kg bodyweight	1 pump per 200g bodyweight (0.2ml) 5 pumps per kg bodyweight (1ml)
Birds	Do not use in animals under 100g bodyweight	200µg/kg bodyweight	1 pump per 200g bodyweight (0.2ml)

Animals under 200g bodyweight should be treated with a single pump actuation.

It is not advisable to treat animals under 100g bodyweight with this product.

(For mammals and birds less than 100g bodyweight use XENO 50–mini)

DO NOT OVERDOSE

(µg = microgram)

For routine prevention of common ectoparasites this dose should be repeated monthly. For the treatment of an existing infestation, repeat doses are required two weeks and four weeks after the initial application.

Special warnings for each target species

Do not overdose. Do not use in animals under 16 weeks of age. Occasionally central nervous system disorders, lethargy and inappetance may be observed (especially in lizards). Avoid spraying in or around the eyes. Local irritation and/or hypersensitivity reactions may occur. Should such a reaction occur, discontinue use and seek veterinary advice. This product is toxic to chelonians. Do not use in tortoises, turtles or related species.

Withdrawal periods

Not applicable. For pet animals only. Do not use on animals intended for human consumption.

Special precautions to be taken by the person administering the product

Wash hands thoroughly with soap and water following treatment.

Pharmaceutical particulars

Major incompatibilities

None known.

Special precautions for storage

Do not store above 25°C. Keep out of direct sunlight.

Keep out of reach of children.

After use, dispose with domestic refuse.

Nature and contents of container

High density polypropylene (HDPE) container containing 50ml with pump aerosol

Marketing authorisation number

WL19886.

Any other information

-

Legal category
SAES
Significant Changes